Journal of Biochemical and Biophysical Methods, 18 (1989) 338–340
Elsevier

Author Index

Balasubramanian, D., see Shobha, J. (18) 287
Balgobin, N., see Remaud, G. (18) 1
Bartosz, G., see Gaczyńska, M. (18) 331
Bause-Niedrig, I., see Sinha, P. (18) 195
Behrendt, G., see Merkel, D. (18) 277
Beile, E., see Berger, R.L. (18) 113
Berger, R.L., Mudd, C.P., Clem, T., Kolobow, T., Beile, E., Simons, P.C., Michel, S. and McClintock, W.
A stopped-flow mixer device for a batch microcalorimeter application to NAD-NADase reaction (18) 113
Bertrand, O., Delfau, M.-H., Garbarz, M., Picat, C., Devaux, I., Dhermy, D., Boivin, P. and Grandchamp, B.
An efficient laboratory made apparatus for DNA amplification (18) 227
Bhattacharyya, B., see Shobha, J. (18) 287
Bianchi-Bosisio, A., see Wajcman, H. (18) 65
Boivin, P., see Bertrand, O. (18) 227
Buck, H.M., see Remaud, G. (18) 1

Centelles, J.J. and Franco, R.
Heterogeneity of the gradients performed by the freeze-thaw method (18) 177
Chai, Y.G., see Singh, B.R. (18) 105
Chan, H.C., Magin, R.L. and Swartz, H.M.
Rapid assessment of liposomal stability in blood by an aqueous nitroxide spin label (18) 271
Chattopadhyaya, J., see Remaud, G. (18) 1
Choi, J., see Singh, B.R. (18) 135
Clem, T., see Berger, R.L. (18) 113
Corbett, J.T.
The scopoletin assay for hydrogen peroxide. A review and a better method (18) 297
Coutelle, C., see Merkel, D. (18) 277
Crouch, R.K., see Mani, V. (18) 91

Delfau, M.-H., see Bertrand, O. (18) 227
Devaux, I., see Bertrand, O. (18) 227
Dhermy, D., see Bertrand, O. (18) 227
Drake, A.F., see Remaud, G. (18) 1

Eijsink, V.G.H., see Van Den Burg, B. (18) 209

Flachmeier, C., see Merkel, D. (18) 277
Franco, R., see Centelles, J.J. (18) 177

Gaczyńska, M. and Bartosz, G.
Standardization of the electron spin resonance method for the proteolytic activity assay (18) 331
Garbarz, M., see Bertrand, O. (18) 227
Gaur, R.K., Paliwal, S., Sharma, P. and Gupta, K.C.
A simple and sensitive spectrophotometric method for the quantitative determination of solid supported amino groups (18) 323
Grandchamp, B., see Bertrand, O. (18) 227
Gruber, G., see Wenisch, E. (18) 309
Gupta, K.C., see Gaur, R.K. (18) 323

Hayashi, H., see Iwaguchi, T. (18) 157
Hellingwerf, K.J., see Westerhoff, H.V. (18) 53
Holma, B., see Thorén, S.A. (18) 149
Hulmes, J.D., see Miedel, M.C. (18) 37
Hunger, H.-D., see Merkel, D. (18) 277

Iwaguchi, T., Shimizu, M. and Hayashi, H.
Analysis of electrophoretic mobility histograms of mouse splenocytes and thymocytes during tumor growth and after combined chemotherapy and immunotherapy (18) 157

Jungbauer, A., see Wenisch, E. (18) 309

Kambayashi, J., see Yokota, M. (18) 97
Katinger, H., see Wenisch, E. (18) 309
Khramtsov, V.V., Panteleev, M.V. and Weiner, L.M.
ESR study of proton transport across phospholipid vesicle membranes (18) 237
Kolobow, T., see Berger, R.L. (18) 113
Koole, L.H., see Remaud, G. (18) 1
Köttgen, E., see Sinha, P. (18) 195
Kreger, S., see Scheerer, J. (18) 221
Kwon, T.-I., see Singh, B.R. (18) 135

0165-022X/88/$03.50 © 1988 Elsevier Science Publishers B.V. (Biomedical Division)

Lawrence, D.S., see Prorok, M. (18) 167
Lichtenstein, A.V., Moiseev, V.L. and Zaboykin, M.M.
Hydroxyapatite-mediated transfer of DNA from diluted solutions to nitrocellulose or nylon membranes (18) 77

Magin, R.L., see Chan, H.C. (18) 271
Mani, V. and Crouch, R.K.
Spin trapping of the superoxide anion: complications in the use of the water-soluble nitroso-aromatic reagent DBNBS (18) 91
Matzenauer, C., see Scheerer, J. (18) 221
McClintock, W., see Berger, R.L. (18) 113
Merkel, D., Schmidt, G., Flachmeier, C., Behrendt, G., Coutelle, C. and Hunger, H.-D.
A new neomycin phosphotransferase II solid phase assay in combination with polyacrylamide sodium dodecylsulphate gel electrophoresis (18) 277
Michel, S., see Berger, R.L. (18) 113
Miedel, M.C., Hulmes, J.D. and Pan, Y.-C.E.
The use of fluorescamine as a detection reagent in protein microcharacterization (18) 37
Moiseev, V.L., see Lichtenstein, A.V. (18) 77
Mori, T., see Yokota, M. (18) 97
Mudd, C.P., see Berger, R.L. (18) 113
Muhn, P., see Scheerer, J. (18) 221
Münzner, P. and Voigt, J.
A convenient procedure for the isolation of intact translatable mRNA by potassium iodide gradient centrifugation (18) 183

Nałęcz, M.J., see Szewczyk, A. (18) 125

Paliwal, S., see Gaur, R.K. (18) 323
Pan, Y.-C.E., see Miedel, M.C. (18) 37
Panteleev, M.V., see Khramtsov, V.V. (18) 237
Picat, C., see Bertrand, O. (18) 227
Prorok, M. and Lawrence, D.S.
Cryopreservation of the cyclic 3′,5′-adenosine monophosphate-dependent protein kinase from bovine cardiac muscle (18) 167

Reiter, M., see Wenisch, E. (18) 309
Remaud, G., Balgobin, N., Sandström, A., Vial, J.-M., Koole, L.H., Buck, H.M., Drake, A.F., Zhou, X.-X. and Chattopadhyaya, J.
Why do all lariat RNA introns have adenosine as the branch-point nucleotide? Conformational study of naturally-occurring branched trinucleotides and its eleven analogues by ^{1}H-, ^{31}P-NMR and CD spectroscopy (18) 1
Righetti, P.G., see Sinha, P. (18) 195
Righetti, P.G., see Wajcman, H. (18) 65
Robertson, D.T., see Singh, B.R. (18) 105

Sakon, M., see Yokota, M. (18) 97
Sandström, A., see Remaud, G. (18) 1
Scheerer, J., Matzenauer, C., Kreger, S. and Muhn, P.
A dot blot assay for sulfated glycosaminoglycans (18) 221
Schill, G.
High performance ion-pair chromatography (18) 249
Schmidt, G., see Merkel, D. (18) 277
Seidel, M., see Sinha, P. (18) 195
Sharma, P., see Gaur, R.K. (18) 323
Shimizu, M., see Iwaguchi, T. (18) 157
Shobha, J., Bhattacharyya, B. and Balasubramanian, D.
Use of micelles in studying drug-binding sites: simulation of the tubulin-bound fluorescence of colchicine (18) 287
Simons, P.C., see Berger, R.L. (18) 113
Singh, B.R., Chai, Y.G., Robertson, D.T. and Song, P.-S.
A photoreversible phytochrome affinity column chromatography for putative phytochrome receptor studies (18) 105
Singh, B.R., Choi, J., Kwon, T.-I. and Song, P.-S.
Use of bilirubin oxidase for probing chromophore topography in tetrapyrrole proteins (18) 135
Sinha, P., Seidel, M., Righetti, P.G., Bause-Niedrig, I. and Köttgen, E.
Use of physiological substrates for zymograms of disaccharidases after separation in immobilized pH gradients (18) 195
Song, P.-S., see Singh, B.R. (18) 105
Song, P.-S., see Singh, B.R. (18) 135
Steindl, F., see Wenisch, E. (18) 309
Stulp, B.K., see Van Den Burg, B. (18) 209
Suurkuusk, J., see Thorén, S.A. (18) 149
Swartz, H.M., see Chan, H.C. (18) 271
Szewczyk, A., Nałęcz, M.J. and Wojtczak, L.
Azido derivatives of dicarboxylic acids for photoaffinity labeling of mitochondrial carriers (18) 125

Tanaka, T., see Yokota, M. (18) 97
Tauer, C., see Wenisch, E. (18) 309
Thorén, S.A., Suurkuusk, J. and Holma, B.
Operation of a multichannel microcalorimetry system in the micro-submicrowatt region: some methodological aspects (18) 149

Tsujinaka, T., see Yokota, M. (18) 97

Van Dam, K., see Westerhoff, H.V. (18) 53
Van Den Burg, B., Eijsink, V.G.H., Stulp, B.K. and Venema, G.
One-step affinity purification of *Bacillus* neutral proteases using Bacitracin-silica (18) 209
Venema, G., see Van Den Burg, B. (18) 209
Vial, J.-M., see Remaud, G. (18) 1
Voigt, J., see Münzner, P. (18) 183

Wajcman, H., Bianchi-Bosisio, A. and Righetti, P.G.
Isoelectric focusing of tryptic peptides from hemoglobin subunits by immobilized pH gradients (18) 65
Weiner, L.M., see Khramtsov, V.V. (18) 237
Wenisch, E., Jungbauer, A., Tauer, C., Reiter, M., Gruber, G., Steindl, F. and Katinger, H.
Isolation of human monoclonal antibody isoproteins by preparative isoelectric focusing in immobilized pH gradients (18) 309
Westerhoff, H.V., Wiechmann, A.H.C.A., Van Dam, K. and Hellingwerf, K.J.
On the evaluation of data from flow-dialysis experiments (18) 53
Wiechmann, A.H.C.A., see Westerhoff, H.V. (18) 53
Wojtczak, L., see Szewczyk, A. (18) 125

Yokota, M., Kambayashi, J., Tanaka, T., Tsujinaka, T., Sakon, M. and Mori, T.
A simple turbidimetric time assay of the endotoxin in plasma (18) 97

Zaboykin, M.M., see Lichtenstein, A.V. (18) 77
Zhou, X.-X., see Remaud, G. (18) 1

Journal of Biochemical and Biophysical Methods, 18 (1989) 341–342
Elsevier

Subject Index

Affinity chromatography, (18) 105; (18) 209
Amino acid analysis, (18) 37
Aminoglycoside antibiotic, (18) 277
Amino group, (18) 323
cAMP-dependent protein kinase, (18) 167
Analysis, (18) 297
Antitumor activity, (18) 157
Autodigestion, (18) 209
Bacilli, (18) 209
Batch microcalorimeter, (18) 113
Bilirubin oxidase, (18) 135
Binding, (18) 105
Binding site feature, (18) 287
Branchpoint nucleotide, (18) 1
Catalase, (18) 91
Circular dichroism, (18) 1
Colchicine emission activation, (18) 287
Collagen, (18) 221
Cryopreservation parameter, (18) 167
Cytochrome *c*, (18) 135
Dicarboxylate carrier, (18) 125
4,4′-Dimethoxytrityl cation, (18) 323
Disaccharidase, (18) 195
DNA, (18) 77
DNA amplification, (18) 227
Dot hybridization, (18) 77
Electron spin resonance, (18) 331
Enantiomeric ion, (18) 249
Endotoxin, (18) 97
Enzyme cryopreservation, (18) 167
ESR, (18) 237; (18) 271
Evaluation algorithm, (18) 53
Fingerprint, (18) 65
Flow dialysis, (18) 53
Fluorescamine detection, (18) 37
Fluorescence, (18) 297
Fluorescence polarization, (18) 287
Freeze-thaw method, (18) 177
Fusion proteins, (18) 277
Glycosaminoglycan, (18) 221
Glycosidase, α-D-, (18) 195
Gradient, (18) 177
Heat exchange rate, (18) 149
Hemoglobin, (18) 65
Heparin, (18) 221
Histone, (18) 221
HPLC, (18) 37
Hydrogen peroxide, (18) 297
Hydroxyapatite, (18) 77
Immobilization dependent fluorescence, (18) 287
Immobilized pH gradient, (18) 65; (18) 195; (18) 309
Immunotherapy, (18) 157
Indirect detection, (18) 249
In vitro translation, (18) 183
Ion-pair distribution, (18) 249
Isoelectric focusing, (18) 65; (18) 195; (18) 309
Isoprotein, (18) 309
KI-gradient centrifugation, (18) 183
Krestin, (18) 157
Limulus test, (18) 97
Liposome, (18) 237; (18) 271
Liquid-solid chromatography, (18) 249
Lymphocyte electrophoresis, (18) 157
Maleimide spin label, (18) 331
Maltase-glucoamylase, (18) 195
Membrane-bound protein, (18) 125
Micellar solubilization, (18) 287
Microcalorimetry, (18) 113; (18) 149
Microheterogeneity, (18) 309
Micro-submicrowatt region, (18) 149
Microviscosity, (18) 287
Mitochondria, (18) 125
Mitomycin C, (18) 157
Molecular weight determination, (18) 277
Monoclonal antibody, (18) 309
Morpholinepropanesulfonic acid, (18) 167
NAD, (18) 113
NADase, (18) 113
Neomycin phosphotransferase II, (18) 277
Neutral protease, (18) 209
Nitrocellulose membrane, (18) 77
Nitroxyl radical, (18) 237
NMR spectroscopy, (18) 1
Northern blot analysis, (18) 183
One-step purification, (18) 209
Organic synthesis, (18) 125
Peptide map, (18) 65

0165-022X/88/$03.50 © 1988 Elsevier Science Publishers B.V. (Biomedical Division)

Peridinin-chlorophyll-*a*-protein, (18) 135
pH probe, (18) 237
Phosphorylation, (18) 277
Photoaffinity labeling, (18) 125
Photomorphogenesis, (18) 105
Phycocyanin-645, (18) 135
Phycocyanin, C, (18) 135
Phytochrome, (18) 105; (18) 135
Polarity, (18) 287
Polymerase chain reaction, (18) 227
Polymer supported amine, (18) 323
Proteinase assay, (18) 331
Protein electroblotting, (18) 37
Protein sequencing, (18) 37
Receptor, (18) 105
Reversed-phase HPLC, (18) 65
RNA, (18) 105
RNA isolation, (18) 183
mRNA, (18) 183
Sampling, (18) 53
Scopoletin, (18) 297
SDS gel electrophoresis, (18) 277
Signal intensity, (18) 271
Sodium 3,5-dibromo-4-nitrosobenzene sulfonate, (18) 91
Spectrophotometric method, (18) 323
Spin trap, (18) 91
Stability, (18) 271
Stopped-flow, (18) 113
Sucrase-isomaltase, (18) 195
Sucrose, (18) 177
Superoxide, (18) 91
Superoxide dismutase, (18) 91
Taq polymerase, (18) 227
Tetrapyrrole, (18) 135
Thermal heat flux, (18) 149
TLC, (18) 37
Transmembrane proton transport, (18) 237
Trehalase, (18) 195
Trinucleotide, (18) 1
Tumor-bearing mice, (18) 157
Turbidimetric assay, (18) 97
Zymogram, (18) 195